

NOVEL 2(1H)-QUINAZOLINONE DERIVATIVE

Bibliographische Daten	Beschreibung	Patentansprüche	Mosaik	Originaldokument	INPADOC Rechtsstand

Veröffentlichungsnummer JP57095966

Veröffentlichungsdatum: 1982-06-15

Erfinder: YAMAMOTO MICHIIHIRO; KOSHIBA MASAO; ORI YUKINORI; AONO SHIYUNJI

Anmelder: SUMITOMO CHEMICAL CO

Klassifikation:

- Internationale: C07D239/78; A61K31/505; A61K31/517; A61P3/06; A61P7/02; A61P9/08; A61P37/08; C07D239/82; C07D401/04; C07D401/12; C07D239/00; A61K31/505; A61K31/517; A61P3/00; A61P7/00; A61P9/00; A61P37/00; C07D401/00; (IPC1-7): A61K31/505; C07D239/78; C07D239/82; C07D401/12

- Europäische:

Anmeldenummer: JP19800171521 19801204

Prioritätsnummer(n): JP19800171521 19801204

399ADOC Patentfamilie anzeigen

Liste der Aufkündigungen: 322

Figure 1

Zusammenfassung von JP57095966

NEW MATERIAL:A compound of formula(R1 is H, alkanoyloxy, formyl, carboxyl, etc.; A is 4-20C hydrocarbon residue; Y is O, S sulfinyl or sulfonyl; R2 is H, halogen or alkyl; R3 is aryl or alkyl; R4 is H, alkyl, hydroxyalkyl, etc.; R6 is H, alkyl, cycloalkyl, alkenyl, etc.; Z is O or S). **EXAMPLE:**1-n-Propyl-6-(3-methyl-2-butenyloxy)-4-phenyl-2(1H)-quinazolin one. **USE:**A medicine having an antileptic, vasodilator action, inhibitory action on the aggregation of blood platelets, antiallergic action, etc. **PROCESS:**A compound of formula III (V is functional group) is reacted with a compound of the formula R1-A-W (W is functional group) to form a (thio)ether bond by the reaction of the groups (V) with (W) and give the compound of formula(Y is O or S).

Daten sind von der **esp@cenet** Datenbank verfügbar - Worldwide

